


Definability, automorphisms and enumeration degrees

Mariya I. Soskova¹

Sofia University and UC Berkeley

Logic Colloquium 2014

¹Supported by a Marie Curie International Outgoing Fellowship STRIDE (298471), BNSF Grant No. DMU 03/07/ 12.12.2011 and Sofia University Science Fund project 97/2014. 

Understanding the structure of the Turing degrees

- 1 Understanding the expressive power of the theory of the Turing degrees.
- 2 Understanding the definable relations in the structure of the Turing degrees.
- 3 Understanding the automorphism group of the Turing degrees.

Understanding the structure of the Turing degrees

- 1 Understanding the expressive power of the theory of the Turing degrees.
 - ▶ Simpson (1977) proved: The theory of \mathcal{D}_T is computably isomorphic to the theory of second order arithmetic
- 2 Understanding the definable relations in the structure of the Turing degrees.
- 3 Understanding the automorphism group of the Turing degrees.

Understanding the structure of the Turing degrees

- 1 Understanding the expressive power of the theory of the Turing degrees.
 - ▶ Simpson (1977) proved: The theory of \mathcal{D}_T is computably isomorphic to the theory of second order arithmetic
- 2 Understanding the definable relations in the structure of the Turing degrees.
 - ▶ Slaman and Woodin (1991) conjectured: The definable relations in \mathcal{D}_T are the ones induced by degree invariant relations on sets definable in second order arithmetic.
- 3 Understanding the automorphism group of the Turing degrees.

Understanding the structure of the Turing degrees

- 1 Understanding the expressive power of the theory of the Turing degrees.
 - ▶ Simpson (1977) proved: The theory of \mathcal{D}_T is computably isomorphic to the theory of second order arithmetic
- 2 Understanding the definable relations in the structure of the Turing degrees.
 - ▶ Slaman and Woodin (1991) conjectured: The definable relations in \mathcal{D}_T are the ones induced by degree invariant relations on sets definable in second order arithmetic.
- 3 Understanding the automorphism group of the Turing degrees.
 - ▶ Slaman and Woodin (1991) conjectured: There are no non-trivial automorphisms of \mathcal{D}_T .

Automorphism bases

Definition

Let \mathcal{A} be a structure. A set $B \subseteq |\mathcal{A}|$ is an automorphism base for \mathcal{A} if whenever f and g are automorphisms of \mathcal{A} such that $(\forall x \in B)(f(x) = g(x))$, then $f = g$.

Automorphism bases

Definition

Let \mathcal{A} be a structure. A set $B \subseteq |\mathcal{A}|$ is an automorphism base for \mathcal{A} if whenever f and g are automorphisms of \mathcal{A} such that $(\forall x \in B)(f(x) = g(x))$, then $f = g$.

Equivalently if f is an automorphism of \mathcal{A} and $(\forall x \in B)(f(x) = x)$ then f is the identity.

Automorphism bases

Definition

Let \mathcal{A} be a structure. A set $B \subseteq |\mathcal{A}|$ is an automorphism base for \mathcal{A} if whenever f and g are automorphisms of \mathcal{A} such that $(\forall x \in B)(f(x) = g(x))$, then $f = g$.

Equivalently if f is an automorphism of \mathcal{A} and $(\forall x \in B)(f(x) = x)$ then f is the identity.

Theorem (Slaman and Woodin)

There is an element $\mathbf{g} \leq \mathbf{0}^{(5)}$ such that $\{\mathbf{g}\}$ is an automorphism base for the structure of the Turing degrees \mathcal{D}_T .

Automorphism bases

Definition

Let \mathcal{A} be a structure. A set $B \subseteq |\mathcal{A}|$ is an automorphism base for \mathcal{A} if whenever f and g are automorphisms of \mathcal{A} such that $(\forall x \in B)(f(x) = g(x))$, then $f = g$.

Equivalently if f is an automorphism of \mathcal{A} and $(\forall x \in B)(f(x) = x)$ then f is the identity.

Theorem (Slaman and Woodin)

There is an element $\mathbf{g} \leq \mathbf{0}^{(5)}$ such that $\{\mathbf{g}\}$ is an automorphism base for the structure of the Turing degrees \mathcal{D}_T .

$\text{Aut}(\mathcal{D}_T)$ is countable and every member has an arithmetically definable presentation.

Every relation induced by a degree invariant definable relation in Second order arithmetic is definable with parameters.

Local structures of Turing degrees

Definition

\mathcal{R} is the substructure of the computably enumerable degrees.

Local structures of Turing degrees

Definition

\mathcal{R} is the substructure of the computably enumerable degrees.

$\mathcal{D}_T(\leq \mathbf{0}')$ is the substructure of all degrees that are bounded by $\mathbf{0}'$, the Δ_2^0 Turing degrees.

Local structures of Turing degrees

Definition

\mathcal{R} is the substructure of the computably enumerable degrees.

$\mathcal{D}_T(\leq \mathbf{0}')$ is the substructure of all degrees that are bounded by $\mathbf{0}'$, the Δ_2^0 Turing degrees.

- 1 Shore (1981) proved that the theory of $\mathcal{D}_T(\leq \mathbf{0}')$ is computably isomorphic to the theory of first order arithmetic.

Local structures of Turing degrees

Definition

\mathcal{R} is the substructure of the computably enumerable degrees.

$\mathcal{D}_T(\leq \mathbf{0}')$ is the substructure of all degrees that are bounded by $\mathbf{0}'$, the Δ_2^0 Turing degrees.

- 1 Shore (1981) proved that the theory of $\mathcal{D}_T(\leq \mathbf{0}')$ is computably isomorphic to the theory of first order arithmetic.
- 2 Harrington and Slaman proved that the theory of \mathcal{R} is computably isomorphic to the theory of first order arithmetic.

The local coding theorem

The local coding theorem

Definition

A set of degrees \mathcal{Z} contained in $\mathcal{D}_T(\leq \mathbf{0}')$ is *uniformly low* if it is bounded by a low degree and there is a sequence $\{Z_i\}_{i < \omega}$, representing the degrees in \mathcal{Z} , and a computable function f such that $\{f(i)\}^{\emptyset'}$ is the Turing jump of $\bigoplus_{j < i} Z_j$.

The local coding theorem

Definition

A set of degrees \mathcal{Z} contained in $\mathcal{D}_T(\leq \mathbf{0}')$ is *uniformly low* if it is bounded by a low degree and there is a sequence $\{Z_i\}_{i < \omega}$, representing the degrees in \mathcal{Z} , and a computable function f such that $\{f(i)\}^{\emptyset'}$ is the Turing jump of $\bigoplus_{j < i} Z_j$.

Example: If $\bigoplus_{i < \omega} A_i$ is low then $\mathcal{A} = \{d_T(A_i) \mid i < \omega\}$ is uniformly low.

The local coding theorem

Definition


A set of degrees \mathcal{Z} contained in $\mathcal{D}_T(\leq \mathbf{0}')$ is *uniformly low* if it is bounded by a low degree and there is a sequence $\{Z_i\}_{i < \omega}$, representing the degrees in \mathcal{Z} , and a computable function f such that $\{f(i)\}^{\emptyset'}$ is the Turing jump of $\bigoplus_{j < i} Z_j$.

Example: If $\bigoplus_{i < \omega} A_i$ is low then $\mathcal{A} = \{d_T(A_i) \mid i < \omega\}$ is uniformly low.

Theorem (Slaman and Woodin)


If \mathcal{Z} is a uniformly low subset of $\mathcal{D}_T(\leq \mathbf{0}')$ then \mathcal{Z} is definable from finitely many parameters in $\mathcal{D}_T(\leq \mathbf{0}')$.

Applications of the coding theorem



Using parameters we can code a model of arithmetic $\mathcal{M} = (\mathbb{N}^{\mathcal{M}}, 0^{\mathcal{M}}, s^{\mathcal{M}}, +^{\mathcal{M}}, \times^{\mathcal{M}}, \leq^{\mathcal{M}})$.


Applications of the coding theorem



Using parameters we can code a model of arithmetic $\mathcal{M} = (\mathbb{N}^{\mathcal{M}}, 0^{\mathcal{M}}, s^{\mathcal{M}}, +^{\mathcal{M}}, \times^{\mathcal{M}}, \leq^{\mathcal{M}})$.

- 1 The set $\mathbb{N}^{\mathcal{M}}$ is definable with parameters \vec{p} .


Applications of the coding theorem



Using parameters we can code a model of arithmetic $\mathcal{M} = (\mathbb{N}^{\mathcal{M}}, 0^{\mathcal{M}}, s^{\mathcal{M}}, +^{\mathcal{M}}, \times^{\mathcal{M}}, \leq^{\mathcal{M}})$.

- 1 The set $\mathbb{N}^{\mathcal{M}}$ is definable with parameters \vec{p} .
- 2 The graphs of s , $+$, \times and the relation \leq are definable with parameters \vec{p} .

Applications of the coding theorem



Using parameters we can code a model of arithmetic $\mathcal{M} = (\mathbb{N}^{\mathcal{M}}, 0^{\mathcal{M}}, s^{\mathcal{M}}, +^{\mathcal{M}}, \times^{\mathcal{M}}, \leq^{\mathcal{M}})$.

- 1 The set $\mathbb{N}^{\mathcal{M}}$ is definable with parameters \vec{p} .
- 2 The graphs of s , $+$, \times and the relation \leq are definable with parameters \vec{p} .
- 3 $\mathbb{N} \models \varphi$ iff $\mathcal{D}_T(\leq \mathbf{0}') \models \varphi_T(\vec{p})$

Applications of the coding theorem

If $\mathcal{Z} \subseteq \mathcal{D}_T(\leq \mathbf{0}')$ is uniformly low and represented by the sequence $\{Z_i\}_{i < \omega}$ then there are parameters that code a model of arithmetic \mathcal{M} and a function $\varphi : \mathbb{N}^{\mathcal{M}} \rightarrow \mathcal{D}_T(\leq \mathbf{0}')$ such that $\varphi(i^{\mathcal{M}}) = d_T(Z_i)$.

Applications of the coding theorem

If $\mathcal{Z} \subseteq \mathcal{D}_T(\leq \mathbf{0}')$ is uniformly low and represented by the sequence $\{Z_i\}_{i < \omega}$ then there are parameters that code a model of arithmetic \mathcal{M} and a function $\varphi : \mathbb{N}^{\mathcal{M}} \rightarrow \mathcal{D}_T(\leq \mathbf{0}')$ such that $\varphi(i^{\mathcal{M}}) = d_T(Z_i)$.

We call such a function *an indexing* of \mathcal{Z} .

Applications of the coding theorem


If $\mathcal{Z} \subseteq \mathcal{D}_T(\leq \mathbf{0}')$ is uniformly low and represented by the sequence $\{Z_i\}_{i < \omega}$ then there are parameters that code a model of arithmetic \mathcal{M} and a function $\varphi : \mathbb{N}^{\mathcal{M}} \rightarrow \mathcal{D}_T(\leq \mathbf{0}')$ such that $\varphi(i^{\mathcal{M}}) = d_T(Z_i)$.

We call such a function *an indexing* of \mathcal{Z} .

Theorem (Slaman and Woodin)

There are finitely many Δ_2^0 parameters which code a model of arithmetic \mathcal{M} and an indexing of the c.e. degrees: a function $\psi : \mathbb{N}^{\mathcal{M}} \rightarrow \mathcal{D}_T(\leq \mathbf{0}')$ such that $\psi(e^{\mathcal{M}}) = d_T(W_e)$.

An indexing of the c.e. degrees



The Goal

The Goal

Extend this result to an indexing φ of the Δ_2^0 Turing degrees.


We will call e an index for a Δ_2^0 set X if $\{e\}^{\emptyset'}$ is the characteristic function of X .

The Goal

The Goal

Extend this result to an indexing φ of the Δ_2^0 Turing degrees.

We will call e an index for a Δ_2^0 set X if $\{e\}^{\emptyset'}$ is the characteristic function of X .




The Goal

The Goal

Extend this result to an indexing φ of the Δ_2^0 Turing degrees.

We will call e an index for a Δ_2^0 set X if $\{e\}^{\emptyset'}$ is the characteristic function of X .



The Goal


The Goal

Extend this result to an indexing φ of the Δ_2^0 Turing degrees.

We will call e an index for a Δ_2^0 set X if $\{e\}^{\emptyset'}$ is the characteristic function of X .

Idea: We can use a further uniformly low set

$$\mathcal{Z} = \{d_T(Z_i) \mid i < \omega\}.$$



The Goal


The Goal

Extend this result to an indexing φ of the Δ_2^0 Turing degrees.

We will call e an index for a Δ_2^0 set X if $\{e\}^{\emptyset'}$ is the characteristic function of X .

Idea: We can use a further uniformly low set

$$\mathcal{Z} = \{d_T(Z_i) \mid i < \omega\}.$$



Biinterpretability with parameters

Theorem (Slaman, S)

There are finitely many Δ_2^0 parameters that code a model of arithmetic \mathcal{M} and an indexing of the Δ_2^0 degrees.

Biinterpretability with parameters

Theorem (Slaman, S)

There are finitely many Δ_2^0 parameters that code a model of arithmetic \mathcal{M} and an indexing of the Δ_2^0 degrees.

Proof flavour:

- 1 A Δ_2^0 degree can be defined from four low degrees using meet and join.

Biinterpretability with parameters

Theorem (Slaman, S)

There are finitely many Δ_2^0 parameters that code a model of arithmetic \mathcal{M} and an indexing of the Δ_2^0 degrees.

Proof flavour:

- 1 A Δ_2^0 degree can be defined from four low degrees using meet and join.
- 2 There exists a uniformly low set of Turing degrees \mathcal{Z} , such that every low Turing degree \mathbf{x} is uniquely positioned with respect to the c.e. degrees and the elements of \mathcal{Z} .

Biinterpretability with parameters

Theorem (Slaman, S)

There are finitely many Δ_2^0 parameters that code a model of arithmetic \mathcal{M} and an indexing of the Δ_2^0 degrees.

Proof flavour:

- 1 A Δ_2^0 degree can be defined from four low degrees using meet and join.
- 2 There exists a uniformly low set of Turing degrees \mathcal{Z} , such that every low Turing degree \mathbf{x} is uniquely positioned with respect to the c.e. degrees and the elements of \mathcal{Z} .

If $\mathbf{x}, \mathbf{y} \leq \mathbf{0}'$, $\mathbf{x}' = \mathbf{0}'$ and $\mathbf{y} \not\leq \mathbf{x}$ then there are $\mathbf{g}_i \leq \mathbf{0}'$, c.e. degrees \mathbf{a}_i and Δ_2^0 degrees $\mathbf{c}_i, \mathbf{b}_i \in \mathcal{Z}$ for $i = 1, 2$ such that:

- 1 \mathbf{g}_i is the least element below \mathbf{a}_i which joins \mathbf{b}_i above \mathbf{c}_i .
- 2 $\mathbf{x} \leq \mathbf{g}_1 \vee \mathbf{g}_2$.
- 3 $\mathbf{y} \not\leq \mathbf{g}_1 \vee \mathbf{g}_2$.

Applications

Theorem (Slaman, S)

- 1 $\mathcal{D}_T(\leq \mathbf{0}')$ has a finite automorphism base.

Applications

Theorem (Slaman, S)

- 1 $\mathcal{D}_T(\leq \mathbf{0}')$ has a finite automorphism base.
- 2 The automorphism group of $\mathcal{D}_T(\leq \mathbf{0}')$ is countable.

Applications

Theorem (Slaman, S)

- 1 $\mathcal{D}_T(\leq \mathbf{0}')$ has a finite automorphism base.
- 2 The automorphism group of $\mathcal{D}_T(\leq \mathbf{0}')$ is countable.
- 3 Every automorphism of $\mathcal{D}_T(\leq \mathbf{0}')$ has an arithmetic presentation.

Applications

Theorem (Slaman, S)

- 1 $\mathcal{D}_T(\leq \mathbf{0}')$ has a finite automorphism base.
- 2 The automorphism group of $\mathcal{D}_T(\leq \mathbf{0}')$ is countable.
- 3 Every automorphism of $\mathcal{D}_T(\leq \mathbf{0}')$ has an arithmetic presentation.
- 4 Every relation $R \subseteq \mathcal{D}_T(\leq \mathbf{0}')$ induced by an arithmetically definable degree invariant relation is definable with finitely many Δ_2^0 parameters.

Applications

Theorem (Slaman, S)

- 1 $\mathcal{D}_T(\leq \mathbf{0}')$ has a finite automorphism base.
- 2 The automorphism group of $\mathcal{D}_T(\leq \mathbf{0}')$ is countable.
- 3 Every automorphism of $\mathcal{D}_T(\leq \mathbf{0}')$ has an arithmetic presentation.
- 4 Every relation $R \subseteq \mathcal{D}_T(\leq \mathbf{0}')$ induced by an arithmetically definable degree invariant relation is definable with finitely many Δ_2^0 parameters.
- 5 $\mathcal{D}_T(\leq \mathbf{0}')$ is rigid if and only if $\mathcal{D}_T(\leq \mathbf{0}')$ is biinterpretable with first order arithmetic.

Enumeration reducibility

Reducibility	Oracle set B	Reduced set A

Enumeration reducibility

Reducibility	Oracle set B	Reduced set A
$A \leq_T B$	Complete information	Complete information

Enumeration reducibility

Reducibility	Oracle set B	Reduced set A
$A \leq_T B$	Complete information	Complete information
A c.e. in B	Complete information	Positive information

Enumeration reducibility

Reducibility	Oracle set B	Reduced set A
$A \leq_T B$	Complete information	Complete information
A c.e. in B	Complete information	Positive information
$A \leq_e B$	Positive information	Positive information

Enumeration reducibility

Reducibility	Oracle set B	Reduced set A
$A \leq_T B$	Complete information	Complete information
A c.e. in B	Complete information	Positive information
$A \leq_e B$	Positive information	Positive information

Definition (Friedberg, Rogers (59))

$A \leq_e B$ if there is a c.e. set W , such that

$$A = W(B) = \{x \mid \exists D(\langle x, D \rangle \in W \ \& \ D \subseteq B)\}.$$

The structure of the enumeration degrees \mathcal{D}_e is an upper semi-lattice with least element and jump operation.

Understanding the structure of the enumeration degrees

Theorem (Slaman, Woodin)

The first order theory of \mathcal{D}_e is computably isomorphic to the theory of Second order arithmetic.

Understanding the structure of the enumeration degrees

Theorem (Slaman, Woodin)

The first order theory of \mathcal{D}_e is computably isomorphic to the theory of Second order arithmetic.

Theorem (S)

The automorphism group of \mathcal{D}_e has the same properties as the automorphism group of \mathcal{D}_T .

What connects \mathcal{D}_T and \mathcal{D}_e

Proposition

$A \leq_T B \Leftrightarrow A \oplus \bar{A}$ is c.e. in $B \Leftrightarrow A \oplus \bar{A} \leq_e B \oplus \bar{B}$.

What connects \mathcal{D}_T and \mathcal{D}_e

Proposition

$$A \leq_T B \Leftrightarrow A \oplus \bar{A} \text{ is c.e. in } B \Leftrightarrow A \oplus \bar{A} \leq_e B \oplus \bar{B}.$$

A set A is *total* if $A \equiv_e A \oplus \bar{A}$. An enumeration degree is *total* if it contains a total set. The set of total degrees is denoted by \mathcal{TOT} .

What connects \mathcal{D}_T and \mathcal{D}_e

Proposition

$$A \leq_T B \Leftrightarrow A \oplus \bar{A} \text{ is c.e. in } B \Leftrightarrow A \oplus \bar{A} \leq_e B \oplus \bar{B}.$$

A set A is *total* if $A \equiv_e A \oplus \bar{A}$. An enumeration degree is *total* if it contains a total set. The set of total degrees is denoted by \mathcal{TOT} .

Example: If f is a total function then G_f is a total set.

What connects \mathcal{D}_T and \mathcal{D}_e

Proposition

$$A \leq_T B \Leftrightarrow A \oplus \bar{A} \text{ is c.e. in } B \Leftrightarrow A \oplus \bar{A} \leq_e B \oplus \bar{B}.$$

A set A is *total* if $A \equiv_e A \oplus \bar{A}$. An enumeration degree is *total* if it contains a total set. The set of total degrees is denoted by \mathcal{TOT} .

Example: If f is a total function then G_f is a total set. The enumeration jump of very set is total.

What connects \mathcal{D}_T and \mathcal{D}_e

Proposition

$$A \leq_T B \Leftrightarrow A \oplus \bar{A} \text{ is c.e. in } B \Leftrightarrow A \oplus \bar{A} \leq_e B \oplus \bar{B}.$$

A set A is *total* if $A \equiv_e A \oplus \bar{A}$. An enumeration degree is *total* if it contains a total set. The set of total degrees is denoted by \mathcal{TOT} .

Example: If f is a total function then G_f is a total set. The enumeration jump of very set is total.

The embedding $\iota : \mathcal{D}_T \rightarrow \mathcal{D}_e$, defined by $\iota(d_T(A)) = d_e(A \oplus \bar{A})$, preserves the order, the least upper bound and the jump operation.

What connects \mathcal{D}_T and \mathcal{D}_e

Proposition

$$A \leq_T B \Leftrightarrow A \oplus \bar{A} \text{ is c.e. in } B \Leftrightarrow A \oplus \bar{A} \leq_e B \oplus \bar{B}.$$

A set A is *total* if $A \equiv_e A \oplus \bar{A}$. An enumeration degree is *total* if it contains a total set. The set of total degrees is denoted by \mathcal{TOT} .

Example: If f is a total function then G_f is a total set. The enumeration jump of very set is total.

The embedding $\iota : \mathcal{D}_T \rightarrow \mathcal{D}_e$, defined by $\iota(d_T(A)) = d_e(A \oplus \bar{A})$, preserves the order, the least upper bound and the jump operation.

$$(\mathcal{D}_T, \leq_T, \vee, ', \mathbf{0}_T) \cong (\mathcal{TOT}, \leq_e, \vee, ', \mathbf{0}_e) \subseteq (\mathcal{D}_e, \leq_e, \vee, ', \mathbf{0}_e)$$

What connects \mathcal{D}_T and \mathcal{D}_e

Proposition

$$A \leq_T B \Leftrightarrow A \oplus \bar{A} \text{ is c.e. in } B \Leftrightarrow A \oplus \bar{A} \leq_e B \oplus \bar{B}.$$

A set A is *total* if $A \equiv_e A \oplus \bar{A}$. An enumeration degree is *total* if it contains a total set. The set of total degrees is denoted by \mathcal{TOT} .

Example: If f is a total function then G_f is a total set. The enumeration jump of very set is total.

The embedding $\iota : \mathcal{D}_T \rightarrow \mathcal{D}_e$, defined by $\iota(d_T(A)) = d_e(A \oplus \bar{A})$, preserves the order, the least upper bound and the jump operation.

$$(\mathcal{D}_T, \leq_T, \vee, ', \mathbf{0}_T) \cong (\mathcal{TOT}, \leq_e, \vee, ', \mathbf{0}_e) \subseteq (\mathcal{D}_e, \leq_e, \vee, ', \mathbf{0}_e)$$

Question (Rogers (67))

Is the set of total enumeration degrees first order definable in \mathcal{D}_e ?

Semi-computable sets

Definition (Jockusch)

A is semi-computable if there is a total computable function s_A , such that $s_A(x, y) \in \{x, y\}$ and if $\{x, y\} \cap A \neq \emptyset$ then $s_A(x, y) \in A$.

Semi-computable sets

Definition (Jockusch)

A is semi-computable if there is a total computable function s_A , such that $s_A(x, y) \in \{x, y\}$ and if $\{x, y\} \cap A \neq \emptyset$ then $s_A(x, y) \in A$.

Example:

- A *left cut* in a computable linear ordering is a semi-computable set.

Semi-computable sets

Definition (Jockusch)

A is semi-computable if there is a total computable function s_A , such that $s_A(x, y) \in \{x, y\}$ and if $\{x, y\} \cap A \neq \emptyset$ then $s_A(x, y) \in A$.

Example:

- A *left cut* in a computable linear ordering is a semi-computable set.
- In particular for any set A consider $L_A = \{\sigma \in 2^{<\omega} \mid \sigma \leq A\}$.

Semi-computable sets

Definition (Jockusch)

A is semi-computable if there is a total computable function s_A , such that $s_A(x, y) \in \{x, y\}$ and if $\{x, y\} \cap A \neq \emptyset$ then $s_A(x, y) \in A$.

Example:

- A *left cut* in a computable linear ordering is a semi-computable set.
- In particular for any set A consider $L_A = \{\sigma \in 2^{<\omega} \mid \sigma \leq A\}$.
- Every nonzero Turing degree contains a semi-computable set that is not c.e. or co-c.e.

Semi-computable sets

Definition (Jockusch)

A is semi-computable if there is a total computable function s_A , such that $s_A(x, y) \in \{x, y\}$ and if $\{x, y\} \cap A \neq \emptyset$ then $s_A(x, y) \in A$.

Example:

- A *left cut* in a computable linear ordering is a semi-computable set.
- In particular for any set A consider $L_A = \{\sigma \in 2^{<\omega} \mid \sigma \leq A\}$.
- Every nonzero Turing degree contains a semi-computable set that is not c.e. or co-c.e.

Theorem (Arslanov, Cooper, Kalimullin)

If A is a semi-computable set then for every X :

$$(d_e(X) \vee d_e(A)) \wedge (d_e(X) \vee d_e(\bar{A})) = d_e(X).$$

Semi-computable sets

Definition (Jockusch)

A is semi-computable if there is a total computable function s_A , such that $s_A(x, y) \in \{x, y\}$ and if $\{x, y\} \cap A \neq \emptyset$ then $s_A(x, y) \in A$.

Example:

- A *left cut* in a computable linear ordering is a semi-computable set.
- In particular for any set A consider $L_A = \{\sigma \in 2^{<\omega} \mid \sigma \leq A\}$.
- Every nonzero Turing degree contains a semi-computable set that is not c.e. or co-c.e.

Theorem (Arslanov, Cooper, Kalimullin)

If A is a semi-computable set then for every X :

$$(d_e(X) \vee d_e(A)) \wedge (d_e(X) \vee d_e(\bar{A})) = d_e(X).$$

- If X is not computable then there is a semi-computable set A with $d_e(X \oplus \bar{X}) = d_e(A) \vee d_e(\bar{A})$.

Kalimullin pairs

Definition (Kalimullin)

A pair of sets A, B are called a \mathcal{K} -pair if there is a c.e. set W , such that $A \times B \subseteq W$ and $\bar{A} \times \bar{B} \subseteq \bar{W}$.

Kalimullin pairs

Definition (Kalimullin)

A pair of sets A, B are called a \mathcal{K} -pair if there is a c.e. set W , such that $A \times B \subseteq W$ and $\overline{A} \times \overline{B} \subseteq \overline{W}$.

Example:

- 1 A trivial example is $\{A, U\}$, where U is c.e: $W = \mathbb{N} \times U$.

Kalimullin pairs

Definition (Kalimullin)

A pair of sets A, B are called a \mathcal{K} -pair if there is a c.e. set W , such that $A \times B \subseteq W$ and $\bar{A} \times \bar{B} \subseteq \bar{W}$.

Example:

- 1 A trivial example is $\{A, U\}$, where U is c.e: $W = \mathbb{N} \times U$.
- 2 If A is a semi-computable set, then $\{A, \bar{A}\}$ is a \mathcal{K} -pair:
 $W = \{(m, n) \mid s_A(m, n) = m\}$.

Kalimullin pairs

Definition (Kalimullin)

A pair of sets A, B are called a \mathcal{K} -pair if there is a c.e. set W , such that $A \times B \subseteq W$ and $\bar{A} \times \bar{B} \subseteq \bar{W}$.

Example:

- 1 A trivial example is $\{A, U\}$, where U is c.e: $W = \mathbb{N} \times U$.
- 2 If A is a semi-computable set, then $\{A, \bar{A}\}$ is a \mathcal{K} -pair:
 $W = \{(m, n) \mid s_A(m, n) = m\}$.

Theorem (Kalimullin)

A pair of sets A, B is a \mathcal{K} -pair if and only if their enumeration degrees \mathbf{a} and \mathbf{b} satisfy:

$$\mathcal{K}(\mathbf{a}, \mathbf{b}) \Leftrightarrow (\forall \mathbf{x} \in \mathcal{D}_e)((\mathbf{a} \vee \mathbf{x}) \wedge (\mathbf{b} \vee \mathbf{x}) = \mathbf{x}).$$

Definability of the enumeration jump

Theorem (Kalimullin)

$\mathbf{0}'_e$ is the largest degree which can be represented as the least upper bound of a triple $\mathbf{a}, \mathbf{b}, \mathbf{c}$, such that $\mathcal{K}(\mathbf{a}, \mathbf{b})$, $\mathcal{K}(\mathbf{b}, \mathbf{c})$ and $\mathcal{K}(\mathbf{c}, \mathbf{a})$.

Definability of the enumeration jump

Theorem (Kalimullin)

$\mathbf{0}'_e$ is the largest degree which can be represented as the least upper bound of a triple $\mathbf{a}, \mathbf{b}, \mathbf{c}$, such that $\mathcal{K}(\mathbf{a}, \mathbf{b})$, $\mathcal{K}(\mathbf{b}, \mathbf{c})$ and $\mathcal{K}(\mathbf{c}, \mathbf{a})$.

Corollary (Kalimullin)

- 1 The enumeration jump is first order definable in \mathcal{D}_e .

Definability of the enumeration jump

Theorem (Kalimullin)

$\mathbf{0}'_e$ is the largest degree which can be represented as the least upper bound of a triple $\mathbf{a}, \mathbf{b}, \mathbf{c}$, such that $\mathcal{K}(\mathbf{a}, \mathbf{b})$, $\mathcal{K}(\mathbf{b}, \mathbf{c})$ and $\mathcal{K}(\mathbf{c}, \mathbf{a})$.

Corollary (Kalimullin)

- 1 The enumeration jump is first order definable in \mathcal{D}_e .
- 2 The set of total enumeration degrees above $\mathbf{0}'_e$ is first order definable in \mathcal{D}_e .

Definability in the local structure of the enumeration degrees

Theorem (Ganchev, S)

The class of \mathcal{K} -pairs below $\mathbf{0}'_e$ is first order definable in $\mathcal{D}_e(\leq \mathbf{0}'_e)$...

Definability in the local structure of the enumeration degrees

Theorem (Ganchev, S)

The class of \mathcal{K} -pairs below $\mathbf{0}'_e$ is first order definable in $\mathcal{D}_e(\leq \mathbf{0}'_e)$...

Theorem (Cai, Lempp, Miller, S)

...by the same formula as in \mathcal{D}_e .

Definability in the local structure of the enumeration degrees

Theorem (Ganchev, S)

The class of \mathcal{K} -pairs below $\mathbf{0}'_e$ is first order definable in $\mathcal{D}_e(\leq \mathbf{0}'_e)$...

Theorem (Cai, Lempp, Miller, S)

...by the same formula as in \mathcal{D}_e .

Theorem (Ganchev, S)

- 1 *The theory of $\mathcal{D}_e(\leq \mathbf{0}'_e)$ is computably isomorphic to the theory of first order arithmetic.*
- 2 *The low enumeration degrees are first order definable in $\mathcal{D}_e(\leq \mathbf{0}'_e)$.*

Maximal \mathcal{K} -pairs

Definition

A \mathcal{K} -pair $\{\mathbf{a}, \mathbf{b}\}$ is maximal if for every \mathcal{K} -pair $\{\mathbf{c}, \mathbf{d}\}$ with $\mathbf{a} \leq \mathbf{c}$ and $\mathbf{b} \leq \mathbf{d}$, we have that $\mathbf{a} = \mathbf{c}$ and $\mathbf{b} = \mathbf{d}$.

Maximal \mathcal{K} -pairs

Definition

A \mathcal{K} -pair $\{\mathbf{a}, \mathbf{b}\}$ is maximal if for every \mathcal{K} -pair $\{\mathbf{c}, \mathbf{d}\}$ with $\mathbf{a} \leq \mathbf{c}$ and $\mathbf{b} \leq \mathbf{d}$, we have that $\mathbf{a} = \mathbf{c}$ and $\mathbf{b} = \mathbf{d}$.

Example: A semi-computable pair is a maximal \mathcal{K} -pair.
Total enumeration degrees are joins of maximal \mathcal{K} -pairs.

Maximal \mathcal{K} -pairs

Definition

A \mathcal{K} -pair $\{\mathbf{a}, \mathbf{b}\}$ is maximal if for every \mathcal{K} -pair $\{\mathbf{c}, \mathbf{d}\}$ with $\mathbf{a} \leq \mathbf{c}$ and $\mathbf{b} \leq \mathbf{d}$, we have that $\mathbf{a} = \mathbf{c}$ and $\mathbf{b} = \mathbf{d}$.

Example: A semi-computable pair is a maximal \mathcal{K} -pair.
Total enumeration degrees are joins of maximal \mathcal{K} -pairs.

Theorem (Ganchev, S)

If $\{A, B\}$ is a nontrivial \mathcal{K} -pair in $\mathcal{D}_e(\leq \mathbf{0}'_e)$ then there is a semi-computable set C , such that $A \leq_e C$ and $B \leq_e \overline{C}$.

Maximal \mathcal{K} -pairs

Definition

A \mathcal{K} -pair $\{\mathbf{a}, \mathbf{b}\}$ is maximal if for every \mathcal{K} -pair $\{\mathbf{c}, \mathbf{d}\}$ with $\mathbf{a} \leq \mathbf{c}$ and $\mathbf{b} \leq \mathbf{d}$, we have that $\mathbf{a} = \mathbf{c}$ and $\mathbf{b} = \mathbf{d}$.

Example: A semi-computable pair is a maximal \mathcal{K} -pair.
Total enumeration degrees are joins of maximal \mathcal{K} -pairs.

Theorem (Ganchev, S)

If $\{A, B\}$ is a nontrivial \mathcal{K} -pair in $\mathcal{D}_e(\leq \mathbf{0}'_e)$ then there is a semi-computable set C , such that $A \leq_e C$ and $B \leq_e \overline{C}$.

Corollary

In $\mathcal{D}_e(\leq \mathbf{0}'_e)$ a nonzero degree is total if and only if it is the least upper bound of a maximal \mathcal{K} -pair.

Defining total enumeration degrees in \mathcal{D}_e

Theorem (Cai, Ganchev, Lempp, Miller, S)

If $\{A, B\}$ is a nontrivial \mathcal{K} -pair in \mathcal{D}_e then there is a semi-computable set C , such that $A \leq_e C$ and $B \leq_e \overline{C}$.

Defining total enumeration degrees in \mathcal{D}_e

Theorem (Cai, Ganchev, Lempp, Miller, S)

If $\{A, B\}$ is a nontrivial \mathcal{K} -pair in \mathcal{D}_e then there is a semi-computable set C , such that $A \leq_e C$ and $B \leq_e \overline{C}$.

Proof flavor: Let W be a c.e. set witnessing that a pair of sets $\{A, B\}$ forms a nontrivial \mathcal{K} -pair.

Defining total enumeration degrees in \mathcal{D}_e

Theorem (Cai, Ganchev, Lempp, Miller, S)

If $\{A, B\}$ is a nontrivial \mathcal{K} -pair in \mathcal{D}_e then there is a semi-computable set C , such that $A \leq_e C$ and $B \leq_e \overline{C}$.

Proof flavor: Let W be a c.e. set witnessing that a pair of sets $\{A, B\}$ forms a nontrivial \mathcal{K} -pair.

- 1 The countable component: we use W to construct an effective labeling of the computable linear ordering \mathbb{Q} .

Defining total enumeration degrees in \mathcal{D}_e

Theorem (Cai, Ganchev, Lempp, Miller, S)

If $\{A, B\}$ is a nontrivial \mathcal{K} -pair in \mathcal{D}_e then there is a semi-computable set C , such that $A \leq_e C$ and $B \leq_e \overline{C}$.

Proof flavor: Let W be a c.e. set witnessing that a pair of sets $\{A, B\}$ forms a nontrivial \mathcal{K} -pair.

- 1 The countable component: we use W to construct an effective labeling of the computable linear ordering \mathbb{Q} .
- 2 The uncountable component: C will be a left cut in this ordering.

Defining total enumeration degrees in \mathcal{D}_e

Theorem (Cai, Ganchev, Lempp, Miller, S)

If $\{A, B\}$ is a nontrivial \mathcal{K} -pair in \mathcal{D}_e then there is a semi-computable set C , such that $A \leq_e C$ and $B \leq_e \overline{C}$.

Proof flavor: Let W be a c.e. set witnessing that a pair of sets $\{A, B\}$ forms a nontrivial \mathcal{K} -pair.

- 1 The countable component: we use W to construct an effective labeling of the computable linear ordering \mathbb{Q} .
- 2 The uncountable component: C will be a left cut in this ordering.

Theorem (Cai, Ganchev, Lempp, Miller, S)

The set of total enumeration degrees is first order definable in \mathcal{D}_e .

The relation *c.e. in*

Definition

A Turing degree \mathbf{a} is *c.e. in* a Turing degree \mathbf{x} if some $A \in \mathbf{a}$ is c.e. in some $X \in \mathbf{x}$.

The relation *c.e. in*

Definition

A Turing degree \mathbf{a} is *c.e. in* a Turing degree \mathbf{x} if some $A \in \mathbf{a}$ is c.e. in some $X \in \mathbf{x}$.

Recall that ι is the standard embedding of \mathcal{D}_T into \mathcal{D}_e .

The relation *c.e. in*

Definition

A Turing degree \mathbf{a} is *c.e. in* a Turing degree \mathbf{x} if some $A \in \mathbf{a}$ is c.e. in some $X \in \mathbf{x}$.

Recall that ι is the standard embedding of \mathcal{D}_T into \mathcal{D}_e .

Theorem (Cai, Ganchev, Lempp, Miller, S)

The set $\{\langle \iota(\mathbf{a}), \iota(\mathbf{x}) \rangle \mid \mathbf{a} \text{ is c.e. in } \mathbf{x}\}$ is first order definable in \mathcal{D}_e .

- 1 Ganchev, S had observed that if \mathcal{TOT} is definable by maximal \mathcal{K} -pairs then the image of the relation ‘c.e. in’ is definable for non-c.e. degrees.
- 2 A result by Cai and Shore allowed us to complete this definition.

The total degrees as an automorphism base

Theorem (Selman)

A is enumeration reducible to B if and only if

$$\{\mathbf{x} \in \mathcal{TOT} \mid d_e(A) \leq \mathbf{x}\} \supseteq \{\mathbf{x} \in \mathcal{TOT} \mid d_e(B) \leq \mathbf{x}\}.$$

The total degrees as an automorphism base

Theorem (Selman)

A is enumeration reducible to B if and only if

$$\{\mathbf{x} \in \mathcal{TOT} \mid d_e(A) \leq \mathbf{x}\} \supseteq \{\mathbf{x} \in \mathcal{TOT} \mid d_e(B) \leq \mathbf{x}\}.$$

Corollary

The total enumeration degrees form a definable automorphism base of the enumeration degrees.

The total degrees as an automorphism base

Theorem (Selman)

A is enumeration reducible to B if and only if

$$\{\mathbf{x} \in \mathcal{TOT} \mid d_e(A) \leq \mathbf{x}\} \supseteq \{\mathbf{x} \in \mathcal{TOT} \mid d_e(B) \leq \mathbf{x}\}.$$

Corollary

The total enumeration degrees form a definable automorphism base of the enumeration degrees.

- *If \mathcal{D}_T is rigid then \mathcal{D}_e is rigid.*

The total degrees as an automorphism base

Theorem (Selman)

A is enumeration reducible to B if and only if

$$\{\mathbf{x} \in \mathcal{TOT} \mid d_e(A) \leq \mathbf{x}\} \supseteq \{\mathbf{x} \in \mathcal{TOT} \mid d_e(B) \leq \mathbf{x}\}.$$

Corollary

The total enumeration degrees form a definable automorphism base of the enumeration degrees.

- *If \mathcal{D}_T is rigid then \mathcal{D}_e is rigid.*
- *The automorphism analysis for the enumeration degrees follows.*

The total degrees as an automorphism base

Theorem (Selman)

A is enumeration reducible to B if and only if

$$\{\mathbf{x} \in \mathcal{TOT} \mid d_e(A) \leq \mathbf{x}\} \supseteq \{\mathbf{x} \in \mathcal{TOT} \mid d_e(B) \leq \mathbf{x}\}.$$

Corollary


The total enumeration degrees form a definable automorphism base of the enumeration degrees.

- *If \mathcal{D}_T is rigid then \mathcal{D}_e is rigid.*
- *The automorphism analysis for the enumeration degrees follows.*
- *The total degrees below $\mathbf{0}_e^{(5)}$ are an automorphism base of \mathcal{D}_e .*

Towards a better automorphism base of \mathcal{D}_e

Theorem (Slaman, Woodin)

There are total Δ_2^0 parameters that code a model of arithmetic \mathcal{M} and an indexing of the image of the c.e. Turing degrees.




Towards a better automorphism base of \mathcal{D}_e

Theorem (Slaman, Woodin)

There are total Δ_2^0 parameters that code a model of arithmetic \mathcal{M} and an indexing of the image of the c.e. Turing degrees.

Idea: In the wider context of \mathcal{D}_e we can reach more elements: non-total elements.



Towards a better automorphism base of \mathcal{D}_e

Theorem (Slaman, S)

If \vec{p} defines a model of arithmetic \mathcal{M} and an indexing of the image of the c.e. Turing degrees then \vec{p} defines an indexing of the total Δ_2^0 enumeration degrees.


Proof flavour:

The image of the c.e. degrees

→ The low 3-c.e. e-degrees


→ The low Δ_2^0 e-degrees

→ The total Δ_2^0 e-degrees




Moving outside the local structure

- 1 Extend to an indexing of all total degrees that are “c.e. in ” and above some total Δ_2^0 enumeration degree.
 - ▶ The jump is definable.
 - ▶ The image of the relation “c.e. in ” is definable.
- 2 Relativizing the previous theorem extend to an indexing of $\bigcup_{\mathbf{x} \leq \mathbf{0}'} \iota([\mathbf{x}, \mathbf{x}'])$.




Moving outside the local structure


- Extend to an indexing of all total degrees below $\mathbf{0}_e''$.




And now we iterate




And now we iterate




And now we iterate



And now we iterate



And now we iterate



Theorem (Slaman, S)

Let n be a natural number and \vec{p} be parameters that index the image of the c.e. Turing degrees. There is a definable from \vec{p} indexing of the total Δ_{n+1}^0 degrees.

Consequences

Theorem (Slaman, S)

- 1 *There is a finite automorphism base for the enumeration degrees consisting of total Δ_2^0 enumeration degrees.*

Consequences

Theorem (Slaman, S)

- 1 *There is a finite automorphism base for the enumeration degrees consisting of total Δ_2^0 enumeration degrees.*
- 2 *The image of the c.e. Turing degrees is an automorphism base for \mathcal{D}_e .*

Consequences

Theorem (Slaman, S)

- 1 *There is a finite automorphism base for the enumeration degrees consisting of total Δ_2^0 enumeration degrees.*
- 2 *The image of the c.e. Turing degrees is an automorphism base for \mathcal{D}_e .*
- 3 *If the structure of the c.e. Turing degrees is rigid then so is the structure of the enumeration degrees.*

Consequences

Theorem (Slaman, S)

- 1 *There is a finite automorphism base for the enumeration degrees consisting of total Δ_2^0 enumeration degrees.*
- 2 *The image of the c.e. Turing degrees is an automorphism base for \mathcal{D}_e .*
- 3 *If the structure of the c.e. Turing degrees is rigid then so is the structure of the enumeration degrees.*

Question

- 1 *Can every automorphism of the Turing degrees be extended to an automorphism of the enumeration degrees?*

Consequences

Theorem (Slaman, S)

- 1 *There is a finite automorphism base for the enumeration degrees consisting of total Δ_2^0 enumeration degrees.*
- 2 *The image of the c.e. Turing degrees is an automorphism base for \mathcal{D}_e .*
- 3 *If the structure of the c.e. Turing degrees is rigid then so is the structure of the enumeration degrees.*

Question

- 1 *Can every automorphism of the Turing degrees be extended to an automorphism of the enumeration degrees?*
- 2 *Can we extend automorphisms of the c.e. degrees to automorphisms of \mathcal{D}_T or of \mathcal{D}_e ?*



Thank you!